

Empirical correction of the EPIC effective area based on *NuSTAR* observations

Felix Fürst • 25.04.2022

A dark blue diagonal gradient bar that starts from the bottom left and extends towards the top right, covering the lower half of the slide.

Overview

Basic Problem

Simultaneous observations between *XMM-Newton* and *NuSTAR* often show cross-calibration residuals (slope, off-set and other small features).

Expected outcome

Correction to EPIC-pn ARF to remove/reduce those features.

Approach

Use simultaneous observations of Crab and 3C 273 and test with sample of AGN.

Caveat

Currently no corrections of *absolute flux offset* implemented.

Crab residuals show a “bump”

Stacked residuals of 13 Crab observations between 2013-2021.

Fitted with same photon index for pn and *NuSTAR*.




Figure 2: Stacked EPIC-pn (red) residuals, FPMA (blue), and FPMB (green) residuals of a joint fit to all Crab epochs, requiring the same photon index for all instrument in each epoch. *a)* Without any correction function for EPIC-pn. *b)* After applying the spline correction function.

Crab residuals show a “bump”

Stacked residuals of 13 Crab observations between 2013-2021.

Fitted with same photon index for pn and *NuSTAR*.

Correct these residuals with a simple spline, anchored at 3 and 12keV.





Figure 3: Spline correction function based on modeling of the Crab nebula. Note that the absolute value of the factor is arbitrarily chosen. In red the best-fit is shown, with the $\pm 1\%$ deviation in orange. In gray an ensemble of 100 solutions based on walkers within the 90% contour of an MCMC simulation are shown, with the corresponding 90% quantile of all walkers shown as a blue dashed line.

Updated correction function – shape



Based on Crab simultaneous observations.

Red line is moved so that below 3keV we have no correction.

This is implemented in
XRT3_XAREAEFF_14.CCF

3C 273 and 1ES 0229+200

Table 2: Observation Log for the 3C 273 and 1ES 0229+200 observations.

Date	XMM	ObsID		Exposure [ks]		Parameters	
		NuSTAR	XMM	NuSTAR	Γ	E_{fold} [keV]	
3C 273							
2012-07-16	0414191001	10002020001	17.23	244.00	$1.620^{+0.010}_{-0.009}$	$(1.98^{+0.30}_{-0.23}) \times 10^2$	
2015-07-13	0414191101	10002020003	47.95	49.42	$1.682^{+0.015}_{-0.021}$	$(2.8^{+1.6}_{-1.0}) \times 10^2$	
2016-06-26	0414191201	10202020002	43.84	35.42	$1.526^{+0.012}_{-0.014}$	$(1.47^{+0.25}_{-0.20}) \times 10^2$	
2017-06-26	0414191301	10302020002	42.42	35.40	1.587 ± 0.014	$(1.64^{+0.44}_{-0.30}) \times 10^2$	
2018-07-04	0414191401	10402020006	42.61	40.32	1.634 ± 0.022	$(1.5^{+0.6}_{-0.4}) \times 10^2$	
2019-07-02	0810820101	10502620002	47.29	49.41	$1.675^{+0.020}_{-0.021}$	$(2.7^{+1.7}_{-0.9}) \times 10^2$	
2020-07-06	0810821501	10602606002	47.63	44.02	$1.622^{+0.014}_{-0.017}$	$(1.6^{+0.5}_{-0.4}) \times 10^2$	
1ES 0229+200							
2021-08-08	0810821801	10702609002	59.70	95.16	1.989 ± 0.023	33^{+7}_{-5}	

3C 273 as secon big calibration source, but spectrum much more complicated than that of the Crab.


But spectral fits show same residuals as in the Crab, and are solved by same correction function!

Apply Correction to 3C 273

7 epochs of 3C 273 data between 2012-2020.


Fitted with `cutoffpl + xillver` (`xillver` normalization fixed in all epochs, distant reflection, see Madsen et al. 2015).

Apply correction shape from Crab fitting; fix FPMA constant to 0.866



3C 273 and 1ES 0229+200

ObsID 0414191201 / 10202020002



Test and checks

AGN Small Window mode Test Sample

Table 4: List of observations used to test the new correction function. All EPIC-pn data were taken in Small Window mode. The last two columns give the best-fit χ^2 value for the uncorrected (base) and corrected data, respectively.


Source	XMM ObsID	NuSTAR ObsID	χ^2_{base}	χ^2_{corr}
NGC 4593	0740920401	60001149006	215.63	216.78
3C120	0693781601	60001042002	286.93	300.55
HE 1136-2304	0741260101	80002031003	222.71	215.87
MCG-6-30-15	0693781401	60001047005	257.04	257.32
HE 1143-1810	0795580101	60302002002	252.38	248.32
HE 1143-1810	0795580201	60302002004	238.57	228.79
HE 1143-1810	0795580501	60302002010	243.22	236.48
Fairall9	0741330101	60001130003	357.50	316.04
MCG-6-30-15	0693781201	60001047002	227.76	219.19
MR2251-178	0763920701	60102025006	263.57	252.70
MR2251-178	0763920801	60102025008	263.70	242.44
MR2251-178	0763920601	60102025004	1550.95	234.60
HE 1143-1810	0795580301	60302002006	259.77	256.14
Swift J2127.4+5654	0693781701	60001110002	236.79	230.14
RXS J1131-1231	0820830101	60401001002	230.55	230.79
Swift J2127.4+5654	0693781901	60001110007	291.14	284.86
MCG-6-30-15	0693781301	60001047003	274.36	254.56
IRAS 09149-6206	0830490101	60401020002	190.32	196.22
ESO511-G030	0852010301	60502035006	239.79	232.78
NGC4151	0679780301	60001111005	327.58	308.40
Mrk359	0830550901	60402021004	231.18	228.56
3C382	0790600201	60202015004	285.77	274.95

Sample based on previous work by
Andrea Gokus and Amy Joyce.


All source rigorously checked for pile-up
and variability (requiring strictly
simultaneous GTIs).

Test with AGN/Small Window sample

No correction



With correction



22 observations from Amy's database.

All fitted individually, simultaneous for NuSTAR and XMM.

Allowed for a free cross calibration.

Showing stacked residuals (chi and ratio)


Full Frame mode Test Sample

Target	XMM ObsID	NuSTAR ObsID	χ^2_{base}	χ^2_{corr}
NGC 1313	0693851201	30002035004	247.55	241.06
IRAS 00521-7054	0795630201	60301029004	242.55	242.55
NGC 4579	0790840201	60201051002	237.82	240.79
CEN X-4	0692790201	30001004002	157.64	156.38
HOLMBERG II X-1	0724810301	30001031005	157.97	158.08
NGC 7090 ULX2	0852050201	80501321002	151.11	151.93
M33 FIELD-2	0800350201	50310002003	150.70	150.57
NGC 1313	0693850501	30002035002	172.35	171.54
M33 FIELD-1	0800350101	50310001004	205.71	204.93
RX J0134.2-4258	0841800201	60501005002	149.98	149.42
ELIAS 29	0800030901	30301001004	151.76	150.01
NGC 1313 X-1	0803990101	30302016002	218.58	218.63
NGC7793 P13	0804670301	30302005002	196.21	193.75
MKN 335	0780500301	80201001002	224.87	223.42
NGC 1313 X-1	0803990601	30302016010	223.06	226.62
HESS J1713-381	0790870201	30201031002	171.40	174.49
IRAS 13197-1627	0763220201	60101020002	312.60	314.37
NGC7793 P13	0804670701	30302005004	178.93	174.06
NGC 1313 X-1	0794580601	90201050002	189.45	191.18
NGC 1313 X-1	0742590301	80001032002	184.97	189.07
NUSTARJ150645+0346.2	0795670101	60301023002	212.49	205.56
NGC 1052	0790980101	60201056002	229.61	227.57
NGC 1194	0852200101	60501011002	248.79	251.82
NGC5907 ULX1	0729561301	80001042002	180.99	183.14
ESO 112-G006	0852180101	60561038002	178.58	178.68
CGCG 475-040	0852181001	60561047002	239.28	240.21
NGC 3081	0852180701	60561044002	288.39	282.98
MGC-07-03-00	0852180201	60561039002	198.14	199.47
ESO 426-G002	0852180301	60561040002	229.07	229.41
NGC 5907 ULX-1	0804091101	30302004008	209.10	206.79
ESO 565-G019	0852180601	60561043002	209.15	208.77
NGC 4785	0743010101	60001143002	206.46	208.11
NGC 6552	0852180901	60561046002	206.08	205.99
ESO 116-18	0795680201	60301027002	227.13	222.70
CXOJ022727.5+333443	0784510301	30201003002	191.64	194.58

Long list of diverse source population in Full Frame mode.

All source rigorously checked for pile-up and variability (requiring strictly simultaneous GTIs).

Full frame mode test sample



Conclusion



Agreement of spectral slope is significantly improved

- Corrections remove a “bump” around 8keV
- for lower S/N sources this “bump” is equivalent to a difference in spectral slope
- Corrections to be released with new CCF files but not applied by default in SAS20.0
- To activate, set “applyabsfluxcorr=yes” in ‘arfgen’ (for both pn and MOS)
- Requesting feedback from the community on the effect of these changes.

Appendix




Next source: 1ES 0229+200

As a possible new good calibration source this BL Lac might be perfect:

- Not too bright / not too faint
- Rather hard power law ($\Gamma \sim 1.8$)
- Simple spectrum (power-law or log-parabola)
- Can even be extended down to UV wavelengths!

Existing data not simultaneously (2009 and 2013)





Crab is extended

EPIC-pn is operated in timing mode.

We need to make sure that *NuSTAR* data is extracted from same source region as CCD4 pn footprint on the sky.

Seasonal change between Northern and Southern are of the remnant.



Absolute Flux correction

NuSTAR observed the Crab as “stray light”, i.e., without the optics.

Avoids largest calibration uncertainty.


Gives a “true” answer to the Crab flux and spectrum:

$$\Gamma = 2.106$$

$$\text{Flux} = 3.368 \times 10^{-8} \text{ erg/cm}^2/\text{s} \text{ (3-50 keV)}$$

i.e., ~12% higher than implied from pointed / calibrated observations

(Madsen et al., 2017b)



Madsen et al., 2017b

NuSTAR correction


In *NuSTAR* we can extract a very large region (200"), covering all of the Crab, just like the stray light.

NuSTAR is already calibrated assuming the Crab has $\Gamma = 2.1$, which is very close to the truth.

Main difference: flux!

Correction factor: focused/stray = 0.866

See also Madsen et al. 2017b





Absolute flux calibration

Current situation (before Oct 2021)

EPIC-pn fluxes are typically
10%-15% **lower** than *NuSTAR*
fluxes.

See also IACHEC paper (Madsen et
al., 2017a)




NuSTAR's new effective area


New *NuSTAR* calibration (from October 2021) **reduced** the effective area by ~10%.

This change is based on comparisons with stray light observations of the Crab (absolute Crab flux) and better understanding of the detector.

This moves *NuSTAR* fluxes **further away** from *XMM-Newton*/EPIC-pn!




Calibration source 3C273 and 1ES 0229



We find a cross-calibration constant for EPIC-pn of $CC_{pn} = 0.8198 \pm 0.0021$ corresponding to a $\sim 18\%$ reduction of the effective area.

Test with AGN/Small Window sample



22 observations from Amy's database.

All fitted individually, simultaneous for NuSTAR and XMM.

Cross calibration fixed at 1.

Showing stacked residuals (chi and ratio)